Open problems on Diophantine m-tuples and elliptic curves

Andrej Dujella

May 17, 2024

As an addition to the book $Diophantine\ m$ -tuples and $Elliptic\ Curves$, we give a selection of open problems and challenging questions. Some are notoriously hard problems, like the question of (un)boundness of ranks of elliptic curves over \mathbb{Q} . Nevertheless, we tried to include certain variants or special cases that might be more tractable. The questions are separated with respect to the corresponding chapters, although there are certainly some questions suitable for more than one chapter.

1 Problems for Chapter 1

Problem 1.1. Find all extensions of the rational Diophantine quadruple

$$\left\{\frac{1}{16}, \frac{33}{16}, \frac{17}{4}, \frac{105}{16}\right\}$$

to a rational Diophantine quintuple.

Problem 1.2. Is there any positive integer $k \ge 2$ such that the Diophantine quadruple $\{k-1, k+1, 4k, 16k^3-4k\}$ has more than one extension to a rational Diophantine quintuple?

Stoll (Acta Arith. **190** (2019), 239–261) proved that the extension is unique for k = 2, 3, 4, 5.

Problem 1.3. Let $\{a, b, c\}$ be a D(-1)-triple and d a positive integer such that ad + 1, bd + 1, and cd + 1 are perfect squares. Do all such quadruples satisfy the formula

$$(d+a+b-c)^2 = 4(ab-1)(cd+1)?$$

Problem 1.4. Do there exist distinct positive integers $a_1, a_2, a_3, b_1, b_2, b_3$ with the property that $a_ib_j + 1$ are perfect squares for each $1 \le i, j \le 3$?

Problem 1.5. Is there any triple a_1, a_2, a_3 of distinct positive integers such that $(a_1x + 1)(a_2x + 1)(a_3x + 1)$ is a perfect square for at least six positive integers x?

Problem 1.6. Prove unconditionally that the size of a set of positive integers with the property that the product of any two of its distinct elements is 1 less than a perfect power is bounded by an absolute constant.

Problem 1.7. Let $D_{m,n}(N)$ denote the number of D(n)-m-tuples with elements in the set $\{1, 2, ..., N\}$ for a given positive integer N.

- (a) If n is a perfect square, is there a constant C(n) depending only on n, such that $D_{3,n} \sim C(n)N\log N$? Is it true that $D(n) = D(1) = \frac{3}{\pi^2}$?
- (b) If n is not a perfect square, is there a constant C(n) depending only on n, such that $D_{3,n} \sim C(n)N$?
- (c) If n is a perfect square, what can be said about the asymptotic behaviour of $D_{4,n}$?
- (d) If n is not a perfect square, is it true that $D_{4,n}$ is finite?

Problem 1.8. Characterize elements z of the ring $\mathbb{Z}[\sqrt{-2}]$ for which there exists a D(z)-quadruple.

Problem 1.9. Formulate a conjecture concerning connections between the existence of D(n)-quadruples and representability of the element n as a difference of two squares in an arbitrary commutative ring with unity, which will be in agreement with all known results and examples on that topic.

Problem 1.10. Define an analogue of a D(n)-m-tuple in a non-commutative ring and prove some non-trivial results concerning the existence of D(n)-quadruples, e.g. in the ring of 2×2 matrices with integer entries.

Problem 1.11. Is there any D(1)-quintuple in the ring of integers of an imaginary quadratic field?

Problem 1.12. Is the size of D(1)-tuples in the ring of integers of a real quadratic field bounded by an absolute constant?

Problem 1.13. Let \mathbb{K} be a number field. Is the size of D(1)-tuples in the ring of integers of \mathbb{K} bounded by a constant depending only on the degree $[\mathbb{K} : \mathbb{Q}]$?

Problem 1.14. Find an upper bound for the size of D(n)-tuples in the ring $\mathbb{Z}[x]$ of polynomials with integer coefficients, which depends only on the degree of n. In particular, find an absolute upper bound in the case of cubic polynomials.

Problem 1.15. Is there any strong Diophantine quadruple, i.e. a set of four non-zero rationals $\{a_1, a_2, a_3, a_4\}$ such that $a_i a_j + 1$ are perfect squares for all $1 \le i, j \le 4$?

Problem 2.1. Are there elliptic curves over \mathbb{Q} of arbitrarily large rank?

Problem 2.2. Is there an elliptic curve over \mathbb{Q} with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$ and rank equal to 4?

Problem 2.3. Are there infinitely many elliptic curves over \mathbb{Q} with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$ and rank ≥ 2 ? The same question for torsion groups $\mathbb{Z}/9\mathbb{Z}$, $\mathbb{Z}/10\mathbb{Z}$, and $\mathbb{Z}/12\mathbb{Z}$.

Problem 2.4. Are there infinitely many elliptic curves over \mathbb{Q} with the torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ (respectively $\mathbb{Z}/8\mathbb{Z}$) and rank > 4?

Problem 2.5. Propose a heuristic to predict the distribution of parities of ranks within known families of elliptic curves over \mathbb{Q} with the torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ (respectively $\mathbb{Z}/8\mathbb{Z}$) and rank ≥ 3 .

Problem 2.6. Are there infinitely many elliptic curves over \mathbb{Q} with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ (respectively $\mathbb{Z}/6\mathbb{Z}$) and rank ≥ 6 ?

Problem 2.7. Is there an infinite family of elliptic curves, parametrized by rational points of a genus 1 curve, with torsion group $\mathbb{Z}/4\mathbb{Z}$ and rank ≥ 7 ?

Problem 2.8. Is there an elliptic curve over a quadratic field with torsion group $\mathbb{Z}/13\mathbb{Z}$, $\mathbb{Z}/15\mathbb{Z}$ or $\mathbb{Z}/18\mathbb{Z}$ and rank ≥ 3 ?

Problem 2.9. Is there an elliptic curve over a quartic field with torsion group $\mathbb{Z}/22\mathbb{Z}$ and positive rank?

Solved by Maksym Voznyy on 2024/01/04.

There is a curve with torsion group $\mathbb{Z}/22\mathbb{Z}$ and rank 2 over the quartic field defined by the polynomial $f(x)=x^4+\frac{50}{27}x^3+\frac{281}{243}x^2+\frac{68}{243}x+\frac{2}{81}$. The curve has the equation $y^2-(201204z^3+328374z^2+159732z+8019)xy-(14996874942306z^3+23680026731268z^2+10746821159640z+931263196176)=x^3-(1435244994z^3+2266248132z^2+1028502360z+89124624)x^2$, where z is a root of f.

Problem 3.1. Is there any rational Diophantine sextuple with at least two integer elements?

Problem 3.2. Is there any rational Diophantine septuple?

Problem 3.3. Are there infinitely many almost rational Diophantine septuples, i.e. sets of seven rationals such that all products increased by 1 are squares, with one exception?

Problem 3.4. Give an explicit absolute bound for the size of rational Diophantine tuples (assuming some plausible conjectures).

Problem 3.5. Is there any elliptic curve over \mathbb{Q} induced by a rational Diophantine triple with rank greater than 12?

Problem 3.6. Is there any elliptic curve over $\mathbb{Q}(t)$ induced by a family of rational Diophantine triples with rank greater than 6?

Problem 3.7. Are there infinitely many elliptic curves over \mathbb{Q} with rank 0 induced by rational Diophantine triples with positive elements?

Problem 3.8. Is there any Diophantine triple $\{a, b, c\}$ such that the elliptic curve $y^2 = (ax + 1)(bx + 1)(cx + 1)$ has torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$?

Problem 3.9. Is there any elliptic curve over $\mathbb{Q}(i)$ induced by a rational Diophantine triple with the torsion group $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ and rank greater than 6?

Problem 3.10. Is there any elliptic curve over \mathbb{Q} induced by a rational Diophantine quadruple with rank greater than 10?

Problem 3.11. Is there any rational Diophantine sextuple which contains a strong Diophantine triple?

By a recent paper by Dujella, Kazalicki and Petričević (arXiv: 2403.17959), there are infinitely many rational Diophantine sextuples which contain, e.g., a strong Diophantine pair {30464/2223, 22815/5168}.

Problem 4.1. Let $k \geq 2$ be an integer. Is it true that the Pellian equation

$$x^2 - (k^2 + 1)y^2 = k^2$$

has at most one solution with 0 < y < k - 1 and x > 0?

Problem 4.2. Is there any positive integer $k, k \neq 33539$, such that the Pellian equation

$$x^2 - (k^2 - 1)y^2 = 2 - k^2$$

has more than four fundamental solutions?

Problem 4.3. Are there infinitely many pairs of Diophantine quadruples with the same largest element?

Problem 4.4. Do there exist two Diophantine quadruples $\{a_1, b, c, d\}$ and $\{a_2, b, c, d\}$ such that $a_1 < a_2 < b < c < d$?

Problem 4.5. Is there any Diophantine quadruple $\{a, b, c, d\}$ such that $a \equiv b \equiv c \equiv d \pmod{k}$ with $k \geq 3$?

Problem 4.6. Are all Diophantine quadruples regular?

Problem 4.7. Are all D(4)-quadruples regular?

Problem 4.8. Let $\{a, b, c\}$ be a Diophantine triple. Do all integer points on the elliptic curve $y^2 = (ax+1)(bx+1)(cx+1)$ satisfy the condition that ax+1, bx+1 and cx+1 are all perfect squares?

Problem 4.9. Let $k \ge 3$ be an integer. Is it true that all integer points on the elliptic curve $y^2 = ((k-1)x+1)((k+1)x+1)(4kx+1)$ are given by $(x,y) \in \{(0,\pm 1), (16k^3-4k, \pm (128k^6-112k^4-20k^2-1))\}$?

Problem 4.10. Propose a heuristic to predict the rank distribution within the family of elliptic curves

$$y^{2} = (F_{2k}x + 1)(F_{2k+2}x + 1)(F_{2k+4}x + 1).$$

Problem 5.1. Is there any D(n)-quintuple for n = -3, 3, 5, 8, 12 or 20?

Problem 5.2. Is the set of all integers n, $n \not\equiv 2 \pmod{4}$, for which there are at most two distinct D(n)-quadruples finite?

Problem 5.3. Is there a polynomial Diophantine quadruple with the property D((4k-1)(4k+1))?

Problem 5.4. Let n be a non-zero integer. Prove that the size of a D(n)-tuple is bounded above by a function of the form $c \log \log |n|$ for an absolute constant c.

Problem 5.5. Is there a third power (respectively a fourth power) Diophantine quadruple?

Problem 5.6. Find an absolute upper bound for the size of $D(p^2)$ -tuples, where p is a prime.

Problem 5.7. Is there any rational D(q)-quintuple for q = 1579?

Problem 5.8. Does a rational D(q)-quintuple exist for every rational number q?

Problem 5.9. Are there infinitely many rational D(q)-quintuples for every rational number q?

Problem 5.10. Is there any rational D(q)-sextuple with q which is not a perfect square?

Problem 5.11. Is there any non-zero integer n and a D(n)-sextuple with all odd elements?

Problem 5.12. Are there infinitely many Diophantine triples that are also $D(n_i)$ -triples for $1 < n_2 < n_3 < n_4$?

Problem 5.13. For a positive integer k, denote by $n_1(k)$ the smallest (in the absolute value) non-zero integer for which there exists a triple $\{a, b, c\}$ of non-zero integers and a set N of integers, such that |N| = k, $n_1(k) \in N$, and $\{a, b, c\}$ is a D(n)-triple for all $n \in N$. Is it true that $n_1(5) = 36$? What can be said (assuming some plausible conjectures) about the asymptotic behaviour of $|n_1(k)|$ for large k?

Problem 5.14. Is there a set of four non-zero integers which is a $D(n_1)$ -, a $D(n_2)$ -, and a $D(n_3)$ -quadruple for three distinct non-zero integers n_1 , n_2 , and n_3 ?

Problem 5.15. Is there a set of five non-zero integers which is a $D(n_1)$ -quintuple and a $D(n_2)$ -quintuple for two distinct non-zero integers n_1 and n_2 ?

Problem 5.16. Is there a rational Diophantine quintuple whose elements are all squares?